

On the X-Ray and Mass Distribution in the Merging Galaxy Cluster 1E 0657-56: Ram Pressure-Stripping in Substructures with an NFW Density Profile

Motokazu TAKIZAWA

*Department of Physics, Yamagata University, Kojirakawa-machi 1-4-12, Yamagata 990-8560
takizawa@sci.kj.yamagata-u.ac.jp*

(Received ; accepted)

Abstract

We investigate the X-ray and mass distribution in the merging galaxy cluster 1E 0657-56. We study head-on collisions of two virialized clusters with an NFW density profile in the Λ CDM universe using an N -body + hydrodynamical code. A clear off-set of an X-ray peak from a mass peak, which is like what is reported in Clowe et al. (2004), is first reproduced in the N -body + hydrodynamical simulations. We estimate the ram pressure-stripping conditions of the substructure in mergers of two NFW dark halos using a simple analytical model. We find that the ram pressure dominates the gravity of the substructure when the smaller cluster's mass is less than approximately one tenth of the larger cluster's mass. The characteristic X-ray and mass structures found in 1E 0657-56 suggest that neither the ram pressure nor the gravitational bound force overwhelms the other and that the mass ratio between the progenitors is near the critical value mentioned above.

Key words: galaxies: clusters: general—galaxies: clusters: individual (1E 0657-56)—galaxies: intergalactic medium—gravitational lensing —hydrodynamics

1. Introduction

According to the standard scenario of cosmological structure formation, clusters of galaxies form through mergers of smaller subclusters. In fact, some of them are forming now. Merging clusters are the sites of structure formation in the universe that can be investigated in detail via different types of observations. Mergers play important roles in cluster evolution itself. Cluster mergers cause bulk flow motion, turbulence, shocks, and/or contact discontinuities in the intracluster medium (ICM), which give us clues to investigate various physical processes in the ICM (see Sarazin 2002 for a review). Turbulence and shocks most likely play crucial roles in particle acceleration in the ICM (e.g. Takizawa, Naito 2000; Ohno, Takizawa, Shibata 2002; Takizawa 2002; Fujita, Takizawa, Sarazin 2003; Brunetti et al. 2003). Strong bulk flow motion and violent pressure changes in the ICM during mergers may affect star formation activities of the member galaxies (Fujita et al. 1999; Bekki, Couch 2003). N -body + hydrodynamical numerical simulations have been carried out to study cluster merger physics (e.g. Schindler, Müller 1993; Ishizaka, Mineshige 1996; Roettiger, Loken, Burns 1997; Takizawa 1999, 2000; Ricker, Sarazin 2001; Ritchie, Thomas 2002). Comparison of such numerical simulations with different kinds of observations give us deep insight into the cluster physics.

1E 0657-56 is one of the most well-known examples of a merging cluster. It is the hottest known cluster and has a very powerful radio halos (Liang et al. 2000). Its simple geometry makes this cluster one of the most suitable

case to investigate cluster merger physics. There are two peaks in both the X-ray surface brightness distribution (Markevitch et al. 2002) and galaxy distribution (Barrena et al. 2002), but their positions do not agree with each other. Observations of the line-of-sight velocities of the member galaxies suggest that its collision axis is almost perpendicular to the line-of-sight (Barrena et al. 2002). From X-ray observations, a bow shock and a cold front are found in front of the smaller subcluster, which suggest that its Mach number is 2 – 3 (Markevitch et al. 2002).

Recently, Clowe et al. (2004) investigated the mass distribution in 1E 0657-56 through weak gravitational lensing. They show clear offsets of the mass density peaks from the X-ray peaks, and that the mass distribution is quite similar to the galaxy one. The smaller substructure in mass is ahead of the X-ray one. They claim that this structure occurs because ICM experience ram pressure but dark matter (DM) and galaxies do not. Although the above-mentioned naive ram pressure-stripping scenario seems to be correct, such characteristic off-sets of X-ray peaks to mass peaks have never been reported in the past numerical simulations. In this paper, we show the first results that successfully reproduce such characteristic structures in the N -body + hydrodynamical simulations, and discuss their implications using a simple analytical model.

The rest of this paper is organized as follows. In §2 we describe the adopted numerical method and initial conditions for our simulations. In §3 we present the simulation results. In §4 we show simple analytical estimation for ram pressure-stripping conditions in mergers of two clus-

ters with an NFW density profile. In §5 we summarize the results.

2. The Simulations

2.1. The Numerical Method

In the present study, we consider clusters of galaxies consisting of two components: collisionless particles corresponding to the galaxies and DM, and gas corresponding to the ICM. When calculating gravity, both components are considered, although the former dominates over the latter. Radiative cooling and heat conduction are not included. We use the Roe total variation diminishing (TVD) scheme to solve the hydrodynamical equations for the ICM (see Hirsch 1990). The hydrodynamical part of the code used here is identical with what is used in Takizawa (2005). Gravitational forces are calculated by the Particle-Mesh (PM) method with the standard FFT technique for the isolated boundary conditions (see Hockney, Eastwood 1988). The size of the simulation box and the number of the grid points are $11.8 \text{ Mpc} \times (5.92 \text{ Mpc})^2$ and $256 \times (126)^2$, respectively. The total number of the N -body particles used in the simulations is $256 \times (128)^2$, which is approximately 4.2×10^6 .

2.2. Models and Initial Conditions

We consider mergers of two virialized subclusters with an NFW density profile (Navarro, Frenk, White 1997) in the Λ CDM universe ($\Omega_0 = 0.25$, $\lambda_0 = 0.75$) for DM. DM masses of the larger and smaller subclusters are $1.00 \times 10^{15} M_\odot$ and $6.25 \times 10^{13} M_\odot$, respectively. Thus, the mass ratio is 16:1. The initial density profiles of the ICM are assumed to be those of a beta-model, where the core radius is a half of the scale radius of the DM distribution, and $\beta = 0.6$. The gas mass fraction is set to be 0.1 inside the virial radius of each subcluster. The parameters such as a virial radius r_v , concentration parameter c for each subcluster are summarized in table 1. We calculate these parameters following the method in Appendix of Navarro, Frenk, White (1997). The radial profiles of the ICM pressure is determined so that the ICM is in hydrostatic equilibrium within the cluster potential of the DM and ICM itself. The resultant temperature profiles are similar to those of initial cluster model in Ricker, Sarazin (2001). The velocity distribution of the DM particles is assumed to be an isotropic Maxwellian. The radial profiles of the DM velocity dispersion are calculated from the Jeans equation with spherical symmetry so that the DM particles are in virial equilibrium in the cluster potential. The coordinate system is taken in such a way that the center of mass is at rest at the origin and that the x -axis is along with the collision axis. The centers of the larger and smaller subclusters are initially located in the sides of $x > 0$ and $x < 0$, respectively. The initial distance between the subcluster's centers is 4.93 Mpc. The initial relative velocity is estimated as in §2 of Takizawa (1999). The resultant value is $8.98 \times 10^2 \text{ km s}^{-1}$, which is approximately two thirds of the infall velocity assuming that they were at rest at infinite distance.

Table 1. Parameters for each subcluster

	$M_{DM} (M_\odot)$	$r_v (\text{Mpc})$	c
cluster 1	1.00×10^{15}	1.97	5.66
cluster 2	6.25×10^{13}	0.784	7.56

3. The Simulation Results

Figure 1(a) shows the X-ray surface brightness distribution (colors) and projected total surface mass density (contours) at a time of 0.67 Gyr after the passage of the subcluster through the core of the larger one. A clear off-set of the mass density peak from the X-ray peak is seen for the smaller subcluster remnant. The mass peak and X-ray peak are located at $x \sim 1.5 \text{ Mpc}$ and $x \sim 1.0 \text{ Mpc}$, respectively. This is clearly because the ICM in the smaller subcluster is lagged by the ram pressure. Figure 1(b) shows the X-ray surface brightness distribution (contours) overlaid with the emissivity-weighted temperature distribution (colors) at the same epoch. A weak jump in the X-ray surface brightness distribution at $x \simeq 1.5$ (near the smaller mass peak) is a bow shock, and the emissivity-weighted temperature is higher in the brighter side, and vice versa. A more prominent jump in the X-ray brightness just in front of the smaller X-ray peak is a contact discontinuity, and the emissivity-weighted temperature is lower in the brighter side, and vice versa. Therefore, this will be recognized as a cold front in actual X-ray observations. Figure 2(a) and (b) show the ICM density and pressure profiles along the collision axis ($y = z = 0$) in front of the smaller X-ray peak, respectively. The bow shock is located at $x \simeq 1.55 \text{ Mpc}$, where jumps are clearly seen in both the density and pressure profiles. The contact discontinuity is at $x \simeq 1.1 \text{ Mpc}$, where a jump is seen only in the density profile and the pressure profile does not have any discontinuity there. As for the overall ICM and DM structures of 1E 0657-56 around the west smaller X-ray and mass peak, our results agree qualitatively with the observations.

4. Discussion on the Ram Pressure-Stripping Conditions

Let us discuss the ram pressure-stripping conditions in merger of two clusters with an NFW DM density profile. We consider the merger of two clusters with masses M_1 and M_2 ($M_1 > M_2$), respectively. We concentrate on the physical status of the ICM in the core region of the smaller subcluster. If the gravity on the subcluster's ICM is weaker than the ram pressure force in unit volume, the ICM will be stripped from the substructure potential. This means,

$$\frac{Gm_2\rho_2}{r_2^2} < A(\pi r_2^2 \rho_1 v^2) \left(\frac{4}{3}\pi r_2^3\right)^{-1}, \quad (1)$$

where G is the gravitational constant, and ρ_1 and ρ_2 are the central gas density of the subcluster 1 and 2, respectively. r_2 and m_2 are the scale radius of the DM profile




Fig. 1. (a) Projected total surface mass density (contours) overlaid with X-ray surface brightness distribution (colors) at a time of 0.67 Gyr after the core passage. A clear off-set of the mass density peak from the X-ray peak is seen. (b) X-ray surface brightness distribution (contours) overlaid with emissivity-weighted temperature distribution (colors) at the same epoch.




Fig. 2. Electron number density (a) and pressure (b) profiles along the collision axis ($y = z = 0$) in front of the substructure at the same epoch as in figure 1. A bow shock is located at $x \simeq 1.55$ Mpc, where jumps are clearly seen in both the density and pressure profiles. A contact discontinuity is at $x \simeq 1.1$ Mpc, where a jump is seen only in the density profiles and the pressure profile has no discontinuity there.

and the DM mass inside r_2 for the cluster 2, respectively. Therefore, the relation between m_2 and M_2 is

$$g(M_2) \equiv \frac{m_2}{M_2} = \frac{\ln 2 - 1/2}{\ln(1+c) - c/(1+c)}, \quad (2)$$

where, c is a concentration parameter of an NFW profile, and weakly depends on the halo mass. A is a fudge factor of an order of unity. It is most likely that $A \lesssim 1$ because all of the ram pressure force is not effective in stripping the gas from the substructure. Some might be used on the excitation of small-scale eddies through Kelvin-Helmholtz instability, and some on the adiabatic compression and shock heating of the ICM, and so on. The collision velocity v has an order of

$$v^2 \simeq \frac{2G(M_1 + M_2)}{R_1 + R_2}, \quad (3)$$

where R_1 and R_2 are the virial radii for the cluster 1 and 2, respectively. It is convenient to introduce a new parameter $\alpha \equiv M_2/M_1$. Then, the stripping condition of inequality (1) becomes

$$F(\alpha : M_1) \equiv \alpha^{2/3-w} \frac{1 + \alpha^{1/3}}{1 + \alpha} - \frac{3A}{2g(\alpha M_1)c(\alpha M_1)} < 0 \quad (4)$$

Deriving inequality (4), we use the scaling relation that $R_2/R_1 = \alpha^{1/3}$ and $\rho_2/\rho_1 = \alpha^{-w}$. In the Λ CDM universe ($\Omega_0 = 0.25$, $\lambda_0 = 0.75$), $w \simeq 1/4$ assuming that ρ_1 and




Fig. 3. Function $F(\alpha : M_1)$ defined by equation (4) for $M_1 = 1.0 \times 10^{15} M_\odot$ in the Λ CDM universe ($\Omega_0 = 0.25$, $\lambda_0 = 0.75$). The solid, dashed, and dot-dashed lines represent the cases of $A = 0.6, 0.4$, and 0.2 , respectively. $F < 0$ means that the ram-pressure dominates the gravity in the substructure.

ρ_2 behave like a characteristic density in an NFW profile (Navarro, Frenk, White 1997).

Figure 3 shows the function $F(\alpha : M_1)$ for $M_1 = 1.0 \times 10^{15} M_\odot$ in the Λ CDM universe ($\Omega_0 = 0.25$, $\lambda_0 = 0.75$). The solid, dashed, and dot-dashed lines represent the cases of $A = 0.6, 0.4$, and 0.2 , respectively. In any case, α is less than ~ 0.1 when $F(\alpha) < 0$. This means that the ram pressure-stripping is more effective for smaller subclusters. Although our estimation here is rather crude, it is interesting that this criteria of $\alpha \sim 0.1$ is close to the mass ratio of our simulations where the clear off-set appears. Obviously, such an off-set does not appear at all if the ram pressure-stripping does not work effectively. The X-ray peak will correspond with the mass peak because the ICM behaves like the DM. On the other hand, if the ram pressure overwhelms the gravity, the ICM in the substructure will not be able to penetrate the larger cluster's center and will be repelled. Furthermore, the larger cluster's ICM is so hot that it cannot be bound by the substructure's gravitational potential. Therefore, we will see a mass peak associated with no X-ray peak. Clear off-sets in the simulation and 1E 0657-56 suggest that the parameter α is close to the above-mentioned critical value where neither the ram pressure nor the gravity absolutely dominates the other.

In our simulations, we do not take into account a dense cool core in the central region of a cluster. However, this probably affects the ram pressure stripping conditions. The existence of the cool core changes ρ_1 and ρ_2 in inequality (1). Thus, the scaling relation $\rho_2/\rho_1 = \alpha^{-w}$ could be modified and/or have some dispersion. In case of 1E0657-56, the smaller subcluster certainly has a cool core, but the larger one seems to have none. Therefore, ρ_2/ρ_1 can be larger than when neither subcluster has a

cooling core. This makes more difficult to strip the ICM.

Whereas the smaller X-ray peak in 1E0657-56 is triangle, that in our simulation results is rather round. The absence of a cool core might cause this difference. Another physical process that may play a crucial role is the magnetic field in the ICM. The magnetic field along the boundary layer probably works to maintain the smaller subcluster's gas as a distinct structure after it is stripped off the DM potential through both the magnetic tension and suppression of heat conduction (Asai, Fukuda, Matsumoto 2004). Dynamical motion of the substructure itself possibly produces this kind of magnetic field configurations (Vikhlinin, Markevitch, Murray 2001). Furthermore, temperature gradients in the boundary layer might produce the magnetic field structure through Weibel-type plasma instabilities (Okabe, Hattori 2003). Three-dimensional high-resolution magnetohydrodynamic simulations will be useful in order to investigate detailed evolution.

Please note that an off-set of an X-ray peak to a mass peak is not a structure in dynamical equilibrium but a transient one. A characteristic timescale of the ram pressure-stripping is estimated to be $\sim R_2/v$, which becomes 0.42 Gyr for the model presented in §2. In the simulation, it takes ~ 0.7 Gyr to strip the ICM from the substructure potential. Certainly this is roughly equal to the above-mentioned value.

5. Summary

We investigate the X-ray and mass structures in the merging galaxy cluster 1E 0657-56. We first reproduce a clear off-set of an X-ray peak to a mass peak in N -body + hydrodynamical simulations of mergers of two subclusters with an NFW and beta-model density profiles for DM and ICM, respectively. As for the overall ICM and DM structures of 1E 0657-56 around the west smaller X-ray and mass peak, our simulation results agree qualitatively with the observations. We discuss the ram pressure-stripping conditions in the mergers of two clusters with an NFW DM density profile using a simple analytical model. We find that the ram pressure dominates the gravity of the substructure when the smaller cluster's mass is less than approximately one tenth of the larger cluster's mass. The characteristic X-ray and DM structures found in 1E 0657-56 suggest that the mass ratio between the progenitors is close to the above-mentioned critical value and that neither the ram-pressure force nor the gravity of the substructure overwhelms the other.

The author would like to thank T. Kitayama, M. Hattori and N. Okabe for valuable discussion, and H. Ishikawa for his contribution in the early stage of this work. The author is also grateful to an anonymous referee for his/her useful comments and suggestions which improved the manuscript. Numerical computations were carried out on VPP5000 at the Center for Computational Astrophysics, CfCA, of the National Astronomical Observatory of Japan. M. T. was supported in part by a Grant-in-Aid from the Ministry of Education,

Science, Sports, and Culture of Japan (16740105).

References

- Asai, N., Fukuda, N., & Matsumoto, R. 2004, *ApJ*, 606, L105
 Barrena, R., Biviano, A., Ramella, M., Falco, E. E., & Seitz, S. 2002, *A&A*, 386, 816
 Bekki, K., & Couch, W. J. 2003, *ApJ*, 596, L13
 Brunetti, G., Blasi, P., Cassano, R., & Gabici, S. 2004, *MNRAS*, 350, 117
 Clowe, D., Gonzalez, A., & Markevitch, M. 2004, *ApJ*, 604, 596
 Fujita, Y., Takizawa, M., Nagashima, M., & Enoki, M. 1999, *PASJ*, 51, L1
 Fujita, Y., Takizawa, M., & Sarazin, C. L. 2003, *ApJ*, 584, 190
 Hirsch, C. 1990, *Numerical Computation of Internal and External Flows* (New York: John Wiley & Sons)
 Hockney, R. W., & Eastwood, J. W. 1988, *Computer Simulation Using Particles* (London: Institute of Physics Publishing)
 Ishizaka, C., & Mineshige, S. 1996, *PASJ*, 48, L37
 Liang, H., Hunstead, R. W., Birkinshaw, M., & Andreani, P. 2000, *ApJ*, 544, 686
 Navarro, J. F., Frenk, C. S., & White, S. D. M. 1997, *ApJ*, 490, 493
 Markevitch, M., Gonzalez, A. H., David, L., Vikhlinin, A., Murray, S., Forman, W., Jones, C., & Tucker, W. 2002, *ApJ*, 567, L27
 Ohno, H., Takizawa, M., & Shibata, S. 2002, *ApJ*, 577, 658
 Okabe, N., & Hattori, M. 2003, *ApJ*, 599, 964
 Roettiger, K., Loken, C., & Burns, J. C. 1997, *ApJS*, 109, 307
 Ricker, P. M., & Sarazin, C. L. 2001, *ApJ*, 561, 621
 Ritchie, B. W., & Thomas, P. A. 2002, *MNRAS*, 329, 675
 Sarazin, C. L. 2002, in *Merging Processes in Galaxy Clusters*, ed. L. Feretti, I. M. Gioia, & G. Giovannini (Dordrecht: Kluwer Academic Publishers), 1
 Schindler, S., & Müller, E. 1993, *A&A*, 272, 137
 Takizawa, M. 1999, *ApJ*, 520, 514
 Takizawa, M. 2000, *ApJ*, 532, 183
 Takizawa, M., & Naito, T. 2000, *ApJ*, 535, 586
 Takizawa, M. 2002, *PASJ*, 54, 363
 Takizawa, M., 2005, *ApJ*, 629, 791
 Vikhlinin, A., Markevitch, M., & Murray, S. S. 2001, *ApJ*, 549, L47